

Ring Structure of SUSY * Product and 1/2 SUSY Wess-Zumino Model

Akifumi Sako[†] and Toshiya Suzuki^{*§}

[†] *Department of Mathematics, Faculty of Science and Technology, Keio University
3-14-1 Hiyoshi, Kohoku-ku, Yokohama 223-8522, Japan*

^{*} *Department of Physics, Faculty of Science, Ochanomizu University*

[§] *Institute of Humanities and Sciences, Ochanomizu University
2-1-1 Otsuka, Bunkyo-ku, Tokyo 112-8610, Japan*

E-mail: [†] sako@math.keio.ac.jp

^{*} tsuzuki@phys.ocha.ac.jp

Abstract

There are two types of non(anti-)commutative deformation of D=4, N=1 supersymmetric field theories and D=2, N=2 theories. One is based on the non-supersymmetric star product and the other is based on the supersymmetric star product. These deformations cause partial breaking of supersymmetry in general. In case of supersymmetric star product, the chirality is broken by the effect of the supersymmetric star product, then it is not clear that lagrangian or observables including F-terms preserve part of supersymmetry. In this article, we investigate the ring structure whose product is defined by the supersymmetric star product. We find the ring whose elements correspond to 1/2 SUSY F-terms. Using this, the 1/2 SUSY invariance of the Wess-Zumino model is shown easily and directly.

PACS codes: 11.10.-z, 11.30.Pb

keywords: non(anti-)commutative deformation, supersymmetric * product, ring structure, 1/2 SUSY

Recently, non(anti-)commutative superspaces have attracted much interest [1] - [32]. Today's activities in this area are strongly motivated by several aspects of the superstring theory, or they occur in connections between supersymmetric field theories and supermatrix models [5],[25]-[32]. Independently of above motivations, from the view point of both theoretical physics and mathematics non(anti-)commutative superspaces are quite interesting subjects. As in cases of ordinary bosonic noncommutative field theories, non(anti-)commutative deformations are implemented by star products [33]. Today, it is known that there are two types of deformations which lead us to non(anti-)commutative superspace: (i) one is based on the non-supersymmetric star product defined by using the supersymmetry generator Q ; (ii) the other is on the supersymmetric star product defined in terms of the covariant derivative D [6]. (In this article, the symbol $\star$ denotes the non-SUSY star product and the symbol $*$ denotes the SUSY star product. Concrete definitions of them will appear below.)

For $D = 4, N = 1$ superspace, as well as for $D = 2, N = 2$ one, we introduce (anti-)chiral superfields. In usual (anti-)commutative supersymmetric field theories, using (anti-)chiral superfields we construct a Wess-Zumino model, which is invariant under the supersymmetry transformation, from F-terms and D-terms. In [5], Seiberg considered the deformation of type (i), and showed that the superalgebra is deformed and 1/2 SUSY survive. (Obeying Seiberg [5], we call 1/2 SUSY as a half of supersymmetry generated by either Q or $\bar{Q}$.) By definition of the $\star$ product if A and B are (anti-)chiral superfield $A \star B$ is trivially (anti-)chiral, in short the $\star$ product does not break (anti-)chirality. As a result, the F-term is invariant under the surviving half supersymmetry. After all, the Wess-Zumino lagrangian is invariant under 1/2 SUSY. On the other hand, in [6], Ferrara *et al.* investigated effects of the type (ii) deformation. There, the scenario of survival of supersymmetry is different from the one in the case of type (i). In contrast to the case (i), the superalgebra is not deformed. However the $*$ product does not preserve (anti-)chirality. As a result, the F-terms break fractions of supersymmetry. They performed explicit calculations for Wess-Zumino models with up to degree 5 superpotential and showed that the F-term breaks 1/2 SUSY and the total lagrangian is invariant under the rest of SUSY.

From these explicit examples of Wess-Zumino models preserving the 1/2 SUSY, it is natural to ask whether fractions of the supersymmetry survive for other superpotentials (more than degree 5) and what observables are invariant under the 1/2 SUSY. For Euclidean case we find the 1/2 SUSY of them by using the result of the non-SUSY $\star$ product theory. But it is not direct way and it does not work for Minkowski case. Therefore it is important subject to understand 1/2 SUSY in terms of the SUSY $*$ product. In the SUSY $*$ product theory, the (anti-)chirality is broken by the effect of the SUSY $*$ product deformation. This breaking makes it difficult to see the 1/2 SUSY invariance. The aim of this letter is to solve this problem and to provide a way of understanding 1/2 SUSY in the framework of SUSY $*$ theory.

We constrain our analysis to a rather simple case. We concentrate on models constructed from a single chiral scalar superfield Φ , which carries no flavor nor color. We find a set of superfields constructed from the chiral scalar superfield and its covariant derivatives. This set has two important properties: (a) it is closed under the $*$ product, i.e. it is a ring defined by using the $*$ product; (b) $\int d\theta^2$ of its element

is invariant under the 1/2 SUSY. As an example, we show that $\Phi_*^n = \overbrace{\Phi * \cdots * \Phi}^n$ belongs to the ring. As a consequence, F-terms constructed of Φ_*^n preserve 1/2 SUSY.

In both (i) and (ii) cases, deformation parameters look like breaking explicit Lorenz invariance. In [5], however, it was shown that the deformation of the Wess-Zumino lagrangian is Lorentz invariant, that is, the deformation parameters appear in the lagrangian through Lorentz invariant combinations. We also check this statement in a bit different way from one in [5].

First of all, we present some formulas, which are useful for the $D = 4, N = 1$ superspace calculation and for the $D = 2, N = 2$ one. Spacetime signature is chosen as Minkowski type in the following expression, but most parts of this article are also valid for Euclid space. Only the [Th3] is restricted to the case of Euclid space, to maintain the hermiticity of the lagrangian.

We start with the $D = 4, N = 1$ case. We use conventions of [34].

The covariant derivatives D_α and $\bar{D}_{\dot{\alpha}}$ are defined by

$$\begin{aligned} D_\alpha &= \partial_{\theta^\alpha} + i\sigma_{\alpha\dot{\alpha}}^\mu \bar{\theta}^{\dot{\alpha}} \partial_\mu , \\ \bar{D}_{\dot{\alpha}} &= -\partial_{\bar{\theta}^{\dot{\alpha}}} - i\theta^\alpha \sigma_{\alpha\dot{\alpha}}^\mu \partial_\mu . \end{aligned} \quad (1)$$

They satisfy the following anti-commutation relations:

$$\{D_\alpha, \bar{D}_{\dot{\alpha}}\} = -2i\sigma_{\alpha\dot{\alpha}}^\mu \partial_\mu , \text{ others} = 0 . \quad (2)$$

The supersymmetry generators Q_α and $\bar{Q}_{\dot{\alpha}}$ are constructed so that they anti-commute with the covariant derivatives D_α and $\bar{D}_{\dot{\alpha}}$. Their explicit forms are given as

$$\begin{aligned} Q_\alpha &= \partial_{\theta^\alpha} - i\sigma_{\alpha\dot{\alpha}}^\mu \bar{\theta}^{\dot{\alpha}} \partial_\mu , \\ \bar{Q}_{\dot{\alpha}} &= -\partial_{\bar{\theta}^{\dot{\alpha}}} + i\theta^\alpha \sigma_{\alpha\dot{\alpha}}^\mu \partial_\mu . \end{aligned} \quad (3)$$

Their anti-commutators produce bosonic translation operators ∂_μ :

$$\{Q_\alpha, \bar{Q}_{\dot{\alpha}}\} = 2i\sigma_{\alpha\dot{\alpha}}^\mu \partial_\mu , \text{ others} = 0 . \quad (4)$$

Using the covariant derivatives, we define (anti-)chiral superfields. The chiral superfields are superfields which are constrained by the following condition,

$$\bar{D}_{\dot{\alpha}}\Phi = 0 . \quad (5)$$

In a similar way, we also define the anti-chiral superfields as

$$D_\alpha\bar{\Phi} = 0 . \quad (6)$$

An important property of the (anti-)chiral superfield, which will be used in the following, is that its highest component is invariant under a half of supersymmetry:

$$QD^2\Phi|_{\bar{\theta}=0} = 0 , \quad \bar{Q}\bar{D}^2\bar{\Phi}|_{\theta=0} = 0 , \quad (7)$$

and becomes some total derivative under the rest:

$$\bar{Q}D^2\Phi|_{\bar{\theta}=0} = \text{total derivative} , \quad Q\bar{D}^2\bar{\Phi}|_{\theta=0} = \text{total derivative} . \quad (8)$$

Now we turn to the $D = 2, N = 2$ case. The superspace coordinates are $(z, \bar{z}, \theta_\pm, \bar{\theta}_\pm)$. All of them can be considered as independent variables. The covariant derivatives $D_\pm$ and $\bar{D}_\pm$ are given as

$$\begin{aligned} D_+ &= \partial_{\theta_-} - i\bar{\theta}_- \partial_z , & D_- &= \partial_{\theta_+} - i\bar{\theta}_+ \partial_{\bar{z}} , \\ \bar{D}_+ &= \partial_{\bar{\theta}_-} - i\theta_- \partial_z , & \bar{D}_- &= \partial_{\bar{\theta}_+} - i\theta_+ \partial_{\bar{z}} , \end{aligned} \quad (9)$$

whose anti-commutation relations are

$$\{D_+, \bar{D}_+\} = -2i\partial_z , \quad \{D_-, \bar{D}_-\} = -2i\partial_{\bar{z}} , \quad \text{others} = 0 . \quad (10)$$

The supersymmetry generators $Q_\pm$ and $\bar{Q}_\pm$, and their anti-commutation relations are given as

$$\begin{aligned} Q_+ &= \partial_{\theta_-} + i\bar{\theta}_- \partial_z , & Q_- &= \partial_{\theta_+} + i\bar{\theta}_+ \partial_{\bar{z}} , \\ \bar{Q}_+ &= \partial_{\bar{\theta}_-} + i\theta_- \partial_z , & \bar{Q}_- &= \partial_{\bar{\theta}_+} + i\theta_+ \partial_{\bar{z}} , \end{aligned} \quad (11)$$

and

$$\{Q_+, \bar{Q}_+\} = +2i\partial_z , \quad \{Q_-, \bar{Q}_-\} = +2i\partial_{\bar{z}} , \quad \text{others} = 0 , \quad (12)$$

respectively.

In a similar way of the $D = 4, N = 1$ case, the (anti-)chiral superfields are defined by

$$\bar{D}_\pm \Phi = 0 , \quad (13)$$

and

$$D_\pm \bar{\Phi} = 0 . \quad (14)$$

They satisfy

$$QD^2\Phi|_{\bar{\theta}=0} = 0 , \quad \bar{Q}\bar{D}^2\bar{\Phi}|_{\theta=0} = 0 , \quad (15)$$

and

$$\bar{Q}D^2\Phi|_{\bar{\theta}=0} = \text{total derivative} , \quad Q\bar{D}^2\bar{\Phi}|_{\theta=0} = \text{total derivative} . \quad (16)$$

Let us introduce non(anti-)commutative deformation into superspaces. Because our main purpose is to investigate the ring structure coming from the $*$ product, it is appropriate to follow the procedure given in [6].

Firstly, we introduce the notion of left/right covariant derivatives. The left covariant derivative is identical to the ordinary covariant derivative

$$\vec{D}\Phi = D\Phi . \quad (17)$$

On the other hand, the right covariant derivative is defined through the following relation

$$\Phi \overleftarrow{D} = (-1)^{p_D(p_\Phi+1)} \vec{D}\Phi , \quad (18)$$

where $p_\mathcal{O}$ denotes the parity of $\mathcal{O}$: for odd quantity $p_{\mathcal{O}_{odd}} = 1$ and for even one $p_{\mathcal{O}_{even}} = 0$. The Leibniz rules hold for both :

$$\begin{aligned} \vec{D}(\Phi\Psi) &= \vec{D}(\Phi)\Psi + (-1)^{p_D p_\Phi} \Phi \vec{D}(\Psi) , \\ (\Phi\Psi) \overleftarrow{D} &= \Phi(\Psi) \overleftarrow{D} + (-1)^{p_D p_\Psi} (\Phi) \overleftarrow{D} \Psi . \end{aligned} \quad (19)$$

In terms of the left and right covariant derivatives, we can define the supersymmetric Poisson bracket $\{ , \}_1$

$$\{\Phi, \Psi\}_1 = P^{\mu\nu} \partial_\mu \Phi \partial_\nu \Psi + P^{\alpha\beta} \Phi \overleftarrow{D}_\alpha \vec{D}_\beta \Psi + \partial_\mu \Phi P^{\mu\alpha} \vec{D}_\alpha \Psi + \Phi \overleftarrow{D}_\alpha P^{\alpha\mu} \partial_\mu \Psi , \quad (20)$$

where $P^{\mu\nu}$ are anti-symmetric, $P^{\alpha\beta}$ symmetric, and $P^{\mu\alpha} = -P^{\alpha\mu}$. If we replace D by $\bar{D}$ in (20), we obtain another supersymmetric Poisson bracket $\{ , \}_2$

$$\{\Phi, \Psi\}_2 = P^{\mu\nu} \partial_\mu \Phi \partial_\nu \Psi + P^{\dot{\alpha}\dot{\beta}} \Phi \overleftarrow{\bar{D}}_{\dot{\alpha}} \vec{\bar{D}}_{\dot{\beta}} \Psi + \partial_\mu \Phi P^{\mu\dot{\alpha}} \vec{\bar{D}}_{\dot{\alpha}} \Psi + \Phi \overleftarrow{\bar{D}}_{\dot{\alpha}} P^{\dot{\alpha}\mu} \partial_\mu \Psi . \quad (21)$$

Also, we can construct non-supersymmetric Poisson bracket $\{ , \}_3$ and $\{ , \}_4$, using Q instead of D :

$$\{\Phi, \Psi\}_3 = P^{\mu\nu} \partial_\mu \Phi \partial_\nu \Psi + P^{\alpha\beta} \Phi \overleftarrow{Q}_\alpha \vec{Q}_\beta \Psi + \partial_\mu \Phi P^{\mu\alpha} \vec{Q}_\alpha \Psi + \Phi \overleftarrow{Q}_\alpha P^{\alpha\mu} \partial_\mu \Psi , \quad (22)$$

$$\{\Phi, \Psi\}_4 = P^{\mu\nu} \partial_\mu \Phi \partial_\nu \Psi + P^{\dot{\alpha}\dot{\beta}} \Phi \overleftarrow{\bar{Q}}_{\dot{\alpha}} \vec{\bar{Q}}_{\dot{\beta}} \Psi + \partial_\mu \Phi P^{\mu\dot{\alpha}} \vec{\bar{Q}}_{\dot{\alpha}} \Psi + \Phi \overleftarrow{\bar{Q}}_{\dot{\alpha}} P^{\dot{\alpha}\mu} \partial_\mu \Psi . \quad (23)$$

Now, we define the supersymmetric star product, using the supersymmetric Poisson bracket of type $\{ , \}_1$. The SUSY star product $(*)$ is defined by

$$\Phi * \Psi = e^P(\Psi, \Phi) = \sum_{n=0}^{\infty} \frac{1}{n!} P^n(\Phi, \Psi) , \quad (24)$$

where

$$P^n(\Phi, \Psi) = \sum_{A_1, \dots, A_n; B_1, \dots, B_n} (-1)^{\rho_{A_1, \dots, A_n}^{B_1, \dots, B_n}} \Phi \overleftarrow{D}_{A_1} \dots \overleftarrow{D}_{A_n} P^{A_1 B_1} \dots P^{A_n B_n} \vec{D}_{B_n} \dots \vec{D}_{B_1} \Psi , \quad (25)$$

and

$$\rho_{A_1, \dots, A_n}^{B_1, \dots, B_n} = \sum_{i=1}^{n-1} (p_{A_i} + p_{B_i}) \sum_{j=i+1}^n p_{A_j} . \quad (26)$$

Notice that non(anti-)commutative parameters P^{AB} are covariantly constant, that is, $D_C P^{AB} = 0$. One can show the associativity of the star product

$$\Phi * (\Psi * X) = (\Phi * \Psi) * X . \quad (27)$$

Replacing D by Q in Eq. (25) makes another star product. This kind of star product is called the non-SUSY star product. We use the symbol $\star$ to denote the non-SUSY star product. It is equivalent to the star product investigated in [5].

Let us consider the $D = 4, N = 1$ case. We do not deal with it for general non(anti-)commutative parameter P^{AB} . Instead, we constrain our analysis to a special case:

$$P^{\alpha\beta} \neq 0 , \text{ others} = 0 . \quad (28)$$

With this setting, we obtain

$$\begin{aligned} \Phi * \Psi &= \Phi \cdot \Psi + P^{\alpha\beta} \Phi \overleftarrow{D}_\alpha \overrightarrow{D}_\beta \Psi + \frac{1}{4} \det P \Phi \overleftarrow{D}^2 \overrightarrow{D}^2 \Psi \\ &= \Phi \cdot \Psi + (-)^{(p_\Phi+1)} P^{\alpha\beta} D_\alpha \Phi D_\beta \Psi - \frac{1}{4} \det P D^2 \Phi D^2 \Psi , \end{aligned} \quad (29)$$

where $\overleftarrow{D}^2 = \overleftarrow{D}_\alpha \overleftarrow{D}_\beta \epsilon^{\alpha\beta}$ and $\overrightarrow{D}^2 = \epsilon^{\beta\alpha} \overrightarrow{D}_\alpha \overrightarrow{D}_\beta$. Remark that the P expansion of the $*$ product terminates at finite order due to the fact that $D_\alpha D_\beta D_\gamma = 0$.

In the $D = 2, N = 2$ case, we use the same setting as (28), so we obtain the same P expansion of the $*$ product (29). (Indices $\alpha, \beta, \dots$ take $+$ or $-$.)

Now preparation is finished, so let us start to investigate the ring structure and 1/2 SUSY of F-terms, and to analyze Lorentz invariance of them. In the following of this article, we treat only the case where 1/2 SUSY is generated by Q . (We can discuss the other case, that is, where 1/2 SUSY corresponds to $\bar{Q}$ by a similar way.) We concentrate on simple models which are constructed from a single chiral scalar superfield Φ with no flavor nor color. So the identity $D_\alpha \Phi D_\beta \Phi D_\gamma \Phi = 0$ holds, which makes our analysis easy. Since the structure of the $*$ product is the same for both the $D = 4, N = 1$ case and the $D = 2, N = 2$ case, which is given by (29), the analysis can be performed in a uniform way. Therefore, statements given in the following are valid for both cases.

Let us suppose following three types of set $X = \{D^\alpha \Phi D_\alpha \Phi\}$, $Y = \{\Phi\}$, $Z = \{D^2 \Phi^n | n = 1, 2, \dots\}$, and let x, y and z_i be their elements, i.e. $x \in X, y \in Y$ and $z_i \in Z$, where $i = 1, 2, \dots$. We introduce following three types of polynomial ring

$$\mathbb{R}[X] = \left\{ \sum_k a_k (D^\alpha \Phi D_\alpha \Phi)^k \right\} = \{a_1 + a_2 D^\alpha \Phi D_\alpha \Phi\} , \quad (30)$$

$$\mathbb{R}[Y] = \left\{ \sum_k a_k \Phi^k \right\} , \quad (31)$$

$$\mathbb{R}[Z] = \left\{ \sum_N \sum_{\{k_1, \dots, k_N\}} a_{k_1, \dots, k_N} \prod_i^N (z_i)^{k_i} | N \in \mathbb{Z}_+ \right\} , \quad (32)$$

where $a_i, a_{k_1, \dots, k_N} \in \mathbb{R}$. (One can replace real number field $\mathbb{R}$ by arbitrary field $\mathbb{F}$. This change does not affect validity of following arguments.) The multiplication of these polynomial rings is determined by ordinary multiplication. Let $R_i(X), R_i(Y)$ and $R_i(Z)$ be polynomials belonging to the polynomial rings i.e. for an arbitrary index i , $R_i(X) \in \mathbb{R}[X]$, $R_i(Y) \in \mathbb{R}[Y]$ and $R_i(Z) \in \mathbb{R}[Z]$. Next step, we define some

sets as follows,

$$D_\alpha \mathbb{R}[X] \equiv \{D_\alpha R_i(X) \mid \forall R_i(X) \in \mathbb{R}[X]\} = \{a_i D_\alpha \Phi D^2 \Phi \mid \forall a_i \in \mathbb{R}\}, \quad (33)$$

$$\begin{aligned} D^2 \mathbb{R}[X] &\equiv \{D^2(R_i(X)) \mid \forall R_i(X) \in \mathbb{R}[X]\} = \{a_i D^2 \Phi D^2 \Phi \mid \forall a_i \in \mathbb{R}\} \\ &\subset \mathbb{R}[Z], \end{aligned} \quad (34)$$

$$\begin{aligned} D_\alpha \mathbb{R}[Y] &\equiv \{D_\alpha R_i(Y) \mid \forall R_i(Y) \in \mathbb{R}[Y]\} = \{D_\alpha \Phi \frac{\delta R_i(Y)}{\delta \Phi} \mid \forall R_i(Y) \in \mathbb{R}[Y]\} \\ &\subset \{(D_\alpha \Phi) R_i(Y) \mid \forall R_i(Y) \in \mathbb{R}[Y]\}, \end{aligned} \quad (35)$$

$$\begin{aligned} D^2 \mathbb{R}[Y] &\equiv \{D^2(R_i(Y)) \mid \forall R_i(Y) \in \mathbb{R}[Y]\} \\ &\subset \{\sum (a_{ij} R_i(Y) R_j(Z) + b_{ij} R_i(X) R_j(Y)) \mid a_{ij}, b_{ij} \in \mathbb{R}\}, \end{aligned} \quad (36)$$

$$D_\alpha \mathbb{R}[Z] \equiv \{D_\alpha R_i(Z) \mid \forall R_i(Z) \in \mathbb{R}[Z]\} = \{0\}, \quad (37)$$

$$D^2 \mathbb{R}[Z] \equiv \{D^2 R_i(Z) \mid \forall R_i(Z) \in \mathbb{R}[Z]\} = \{0\}. \quad (38)$$

We define $\mathbb{R}[XYZ]$ by the set of all polynomials that are produced by the elements of X , Y and Z :

$$\begin{aligned} \mathbb{R}[XYZ] &\equiv \left\{ \sum_{ijk} R_i(X) R_j(Y) R_k(Z) \mid R_i(X) \in \mathbb{R}[X], R_j(Y) \in \mathbb{R}[Y], R_k(Z) \in \mathbb{R}[Z] \right\} \\ &= \left\{ (D^\alpha \Phi D_\alpha \Phi) \sum_N \sum a_{k,k_1,\dots,k_N} \Phi^k (D^2 \Phi)^{k_1} \dots (D^2 \Phi^N)^{k_N} \right. \\ &\quad \left. + \sum_N \sum b_{k,k_1,\dots,k_N} \Phi^k (D^2 \Phi)^{k_1} \dots (D^2 \Phi^N)^{k_N} \right\}, \end{aligned} \quad (39)$$

where $a_{k,k_1,\dots,k_N}$ and $b_{k,k_1,\dots,k_N}$ are C-number coefficients.

Note that $\mathbb{R}[XYZ]$ is a polynomial ring produced by the elements of X , Y and Z whose product is defined by ordinary multiplication.

Let us prove the following theorem that is a key to understand the 1/2 SUSY invariance of the Wess-Zumino action.

Theorem 1 (* Ring)

Take $\mathbb{R}[XYZ]$ and $*$ product as above. Then $\mathbb{R}[XYZ]$ is a polynomial ring constructed by the elements of X , Y and Z whose product is defined by $*$ product i.e. if R_1 and R_2 belong to $\mathbb{R}[XYZ]$ then $R_1 \pm R_2 \in \mathbb{R}[XYZ]$ and $R_1 * R_2 \in \mathbb{R}[XYZ]$.

Proof

It is enough for the proof that we show $\mathbb{R}[XYZ]$ is closed under the $*$ product. For an arbitrary element of $\mathbb{R}[XYZ]$, $R_m[XYZ] \equiv \sum_{ijk} R_i(X) R_j(Y) R_k(Z) \in \mathbb{R}[XYZ]$, $\exists R_{m'}[XYZ] \in \mathbb{R}[XYZ]$ that satisfies

$$\begin{aligned} D_\alpha R_m[XYZ] &= \sum_{ijk} \{(D_\alpha R_i(X)) R_j(Y) R_k(Z) + R_i(X) (D_\alpha R_j(Y)) R_k(Z) \\ &\quad + R_i(X) R_j(Y) (D_\alpha R_k(Z))\} \\ &= D_\alpha \Phi R_{m'}[XYZ]. \end{aligned} \quad (40)$$

We use (33) ~ (38) here. Similarly, we find that $D^2 R_m[XYZ]$ belongs to $\mathbb{R}[XYZ]$, i.e.

$$D^2 R_m[XYZ] \in \mathbb{R}[XYZ]. \quad (41)$$

Using these results, one can show that for $\forall R_m[XYZ], R_n[XYZ] \in \mathbb{R}[XYZ]$,

$\exists R_{m'}, R_{n'}, R_p, R_q \in \mathbb{R}[XYZ]$ that satisfy

$$\begin{aligned}
R_m[XYZ] * R_n[XYZ] &= R_m[XYZ] R_n[XYZ] \\
&\quad + P^{\alpha\beta} R_m[XYZ] \overleftarrow{D}_\alpha \overrightarrow{D}_\beta R_n[XYZ] \\
&\quad + \frac{1}{4} \det P R_m[XYZ] \overleftarrow{D}^2 \overrightarrow{D}^2 R_n[XYZ] \\
&= R_m[XYZ] R_n[XYZ] \\
&\quad + P^{\alpha\beta} D_\alpha \Phi D_\beta \Phi R_{m'}[XYZ] R_{n'}[XYZ] \\
&\quad + R_p[XYZ] R_q[XYZ].
\end{aligned} \tag{42}$$

Because of symmetric property of $P^{\alpha\beta}$, the second term vanishes, i.e.

$$P^{\alpha\beta} D_\alpha \Phi D_\beta \Phi R_{m'}[XYZ] R_{n'}[XYZ] = 0, \tag{43}$$

then we can conclude $R_m[XYZ] * R_n[XYZ] \in \mathbb{R}[XYZ]$. ■

Let us see the relation between $\mathbb{R}[XYZ]$ and 1/2 SUSY, here.

Theorem 2 (Ring and 1/2 SUSY)

Take $\mathbb{R}[XYZ]$ as above. If $R_i[XYZ] \in \mathbb{R}[XYZ]$, then $\int d^2\theta R_i[XYZ]$ is invariant under 1/2 SUSY transformation.

Proof

Arbitrary $R_i[XYZ] \in \mathbb{R}[XYZ]$ is expressed as

$$\begin{aligned}
R_i[XYZ] &= \sum a_{nk} (D^\alpha \Phi D_\alpha \Phi) (\Phi)^n R_k[Z] \\
&\quad + \sum b_{nl} (\Phi)^n R_l[Z],
\end{aligned} \tag{44}$$

where $R_k(Z)$ and $R_l(Z)$ are elements of $\mathbb{R}[Z]$. When D^2 operates on the first term of Eq. (44),

$$\begin{aligned}
D^2 \sum a_{nk} (D^\alpha \Phi D_\alpha \Phi) (\Phi)^n R_k[Z] \\
&= - \sum a_{nk} D^2 \Phi D^2 \Phi (\Phi)^n R_k[Z] - \sum a_{nk} n (D^2 \Phi) (D^\alpha \Phi D_\alpha \Phi) \Phi^{n-1} R_k[Z] \\
&= - \sum a_{nk} \frac{1}{n+1} (D^2 \Phi) (D^2 (\Phi)^{n+1}) R_k[Z].
\end{aligned} \tag{45}$$

When D^2 operates on the second term of Eq. (44),

$$D^2 \sum b_{nl} (\Phi)^n R_l[Z] = \sum b_{nl} (D^2 (\Phi)^n) R_l[Z]. \tag{46}$$

Eqs.(45) and (46) show that $D^2 R_i[XYZ] \in \mathbb{R}[Z]$. Then some $R_j[Z] (\in \mathbb{R}[Z])$ exists that satisfies

$$\int d^2\theta R_i[XYZ] = D^2 R_i[XYZ]|_{\bar{\theta}=0} = R_j[Z]|_{\bar{\theta}=0}. \tag{47}$$

Recall that $R_j[Z]$ is some polynomial of D^2 exact terms and D^2 exact terms are invariant under Q (see Eq.(7) or Eq.(15)). Therefore, it is proved that $\int d^2\theta R_i[XYZ]$ is invariant under 1/2 SUSY transformation. ■

Using above theorems, we can easily show that 1/2 SUSY invariance of F-terms.

Theorem 3 (1/2 SUSY invariance of F-terms)

Take $*$ product as above. Let Φ be a chiral superfield. Then, for arbitrary $n \in \mathbb{N}$, $\int d^2\theta(\Phi)_*^n = \int d^2\theta\Phi * \dots * \Phi$ is invariant under 1/2 SUSY transformation.

Proof

Because of the theorem of Ring structure [Th1], $(\Phi)_*^n \in \mathbb{R}[XYZ]$. From [Th2], it is proved that $\int d^2\theta(\Phi)_*^n$ is 1/2SUSY invariant. ■

The F-term 1/2 SUSY of the $*$ product is shown by the F-term 1/2 SUSY of the $\star$ product without the ring $\mathbb{R}[XYZ]$, as follows. When we consider the Euclidean space, we can take $\bar{\theta} = 0$ before $\int d\theta^2$. Therefore replacing the D operators with the Q operators makes no difference in the computation of the F-terms. In short, we can exchange $*$ by $\star$ in F-terms and the F-terms are identical in both cases. Since the $\star$ product does not break chirality, this fact implies the 1/2 SUSY of the $\int d^2\theta(\Phi)_*^n$.¹

In order to see other usefulness of the ring $\mathbb{R}[XYZ]$, let us construct *non-trivial* observables which can have non-zero v.e.v. in 1/2 SUSY invariant phases. Consider observables

$$\int d\theta^2 D^\alpha * \Phi * D_\alpha * \Phi * (\Phi)_*^n = \int d\theta^2 ((D^\alpha * \Phi) * (D_\alpha * \Phi)) * (\Phi)_*^n. \quad (48)$$

Since $D_\alpha * \Phi = D_\alpha \Phi$ and $D^\alpha \Phi * D_\alpha \Phi = D^\alpha \Phi D_\alpha \Phi$, the integrand of (48) is equal to $(D^\alpha \Phi D_\alpha \Phi) * (\Phi)_*^n$. Because $(D^\alpha \Phi D_\alpha \Phi)$ and $(\Phi)_*^n$ belong to $\mathbb{R}[XYZ]$, we conclude $\int d\theta^2 D^\alpha * \Phi * D_\alpha * \Phi * (\Phi)_*^n$ is 1/2 SUSY invariant by [Th1] and [Th2].

In the $\star$ theory, the 1/2 SUSY invariance of $\int d\theta^2 D^\alpha \star \Phi \star D_\alpha \star \Phi \star (\Phi)_*^n$, the counterpart of (48), is not manifest, as they include non-chiral objects. But, by making use of 1/2 SUSY of (48), we conclude that they are still 1/2 SUSY invariant.² Thus the ring $\mathbb{R}[XYZ]$ provides a new method to construct 1/2 SUSY invariant observables which explicitly break (anti-)chirality.

We have studied the relation between the 1/2 SUSY and the $*$ product above. Not only the SUSY but also the Lorentz invariance of the theories becomes nontrivial under the non(anti-)commutative deformation. The following theorem solves this problem.

Theorem 4 (Lorentz invariance)

Let f be some Lorentz invariant superfield. We denote f_*^n as $\overbrace{f * \dots * f}^n$. Then some Lorentz invariant functional $g(f, (Df)^2, D^2 f; \det P)$ exist, where $(Df)^2 = D^\alpha f D_\alpha f$, and it satisfies that

$$f_*^n = f^n + g. \quad (49)$$

Here the noncommutative parameters $P^{\alpha\beta}$ dependence only appear as the $\det P$ dependence. This fact shows that f_*^n is Lorentz invariant.

Proof

We prove this theorem by using mathematical induction as follows.

(i) $n = 2$;

$$f * f = f^2 - \frac{1}{4}(\det P)(D^2 f)^2 \quad (50)$$

(ii) Suppose

$$f_*^n = f^n + g. \quad (51)$$

¹This argument was taught us by the referee of Phys. Lett. B.

²As noted in [13], $Q_\alpha \star \Phi$ is a chiral superfield in the $\star$ theory. Then by using the same argument as above 1/2 SUSY of $\int d\theta^2 D^\alpha \star \Phi \star D_\alpha \star \Phi \star (\Phi)_*^n$ is shown. We thank the referee of Phys. Lett. B for pointing out this.

Using this,

$$f_*^{n+1} = f * f^n + f * g. \quad (52)$$

The first term is rewritten as

$$f * f^n = f^{n+1} - \frac{1}{4}(\det P)(D^2 f)(D^2 f^n). \quad (53)$$

The second term is given as

$$f * g = fg + (-)^{p_f+1} P^{\alpha\beta} D_\alpha f D_\beta g - \frac{1}{4}(\det P)(D^2 f)(D^2 g). \quad (54)$$

We can show that the second term of Eq.(54) vanishes, as follows.

$$\begin{aligned} D_\alpha g(f, (Df)^2, D^2 f) &= (D_\alpha f) \frac{\partial g}{\partial f} + (D_\alpha (Df)^2) \frac{\partial g}{\partial ((Df)^2)} + (D_\alpha (D^2 f)) \frac{\partial g}{\partial (D^2 f)} \\ &= (D_\alpha f) \frac{\partial g}{\partial f} - \sum_\beta \epsilon^{\alpha\beta} (D_\alpha D_\beta f) D_\alpha f \frac{\partial g}{\partial ((Df)^2)}. \end{aligned} \quad (55)$$

Here we do not sum over the index α . Using Eq.(55) and symmetric nature of $P^{\alpha\beta}$, the second term of Eq.(54) vanishes ;

$$\begin{aligned} &\sum_{\alpha\beta} P^{\alpha\beta} D_\alpha f D_\beta g \\ &= \sum_{\alpha\beta\gamma} \left\{ \frac{1}{2} P^{\alpha\beta} D_\alpha f D_\beta f \frac{\partial g}{\partial f} - \frac{1}{2} P^{\alpha\beta} (D_\alpha f) \epsilon^{\beta\gamma} (D_\beta D_\gamma f) D_\beta f \frac{\partial g}{\partial ((Df)^2)} \right\} = 0. \end{aligned} \quad (56)$$

Therefore,

$$f_*^{n+1} = f^{n+1} - \frac{1}{4}(\det P)(D^2 f)(D^2 f^n) + fg - \frac{1}{4}(\det P)(D^2 f)(D^2 g) = f^{n+1} + g', \quad (57)$$

which is what we want. Here we denote g' as $-\frac{1}{4}(\det P)(D^2 f)(D^2 f^n) + fg - \frac{1}{4}(\det P)(D^2 f)(D^2 g)$. By the principle of mathematical induction, for all $n \geq 2$ (and $n = 1$ that is trivial case), f_*^n are Lorentz invariant.

■

Note that only the symmetric property of the noncommutative parameter $P^{\alpha\beta}$ is used in this proof. So, we can show the Lorentz invariance for other star products that is defined by other Poisson brackets like the non-SUSY Poisson bracket [5] .

It is worth while to comment here about the ring structure of Lorentz invariant functionals. Let $\mathbb{R}_L[f, (Df)^2, D^2 f; \det P]$ be a set of all polynomials of f , $(Df)^2$ and $D^2 f$, with additional dependence of $\det P$. Since non(anti-)commutative parameter $P^{\alpha\beta}$ appears in elements of $\mathbb{R}_L[f, (Df)^2, D^2 f; \det P]$ only through $\det P$, the elements are Lorentz invariant. We can show that $\mathbb{R}_L[f, (Df)^2, D^2 f; \det P]$ is a ring whose product is defined by the $*$ product, by a similar way to the proof of [Th 1].

Wess-Zumino models contain not only F-terms but also D-terms. In general, D-terms are deformed by the $*$ product, too. However, we can show the Lorentz invariance and the SUSY invariance of the D-terms as follows. Let Kähler potential $K(\Phi, \bar{\Phi})_*$ be defined as formal power series of Φ and $\bar{\Phi}$, where

the multiplication is defined by the SUSY $*$ product. Using the fact that $\Phi * \bar{\Phi} = \Phi \cdot \bar{\Phi}$ and [Th4], the Kähler potential is rewritten as

$$\begin{aligned} K(\Phi, \bar{\Phi})_* &= \sum_{ij} c_{ij} \Phi_*^i \cdot \bar{\Phi}^j \\ &= \sum_{ij} c_{ij} \{ \Phi^i + g_i(\Phi, (D\Phi)^2, D^2\Phi^m; \det P) \} \cdot \bar{\Phi}^j, \end{aligned} \quad (58)$$

where $g_i(\Phi, (D\Phi)^2, D^2\Phi^m; \det P)$ are Lorentz invariant functionals. Eq. (58) shows the Lorentz invariance of the D-term. In addition, $\int d^4\theta K(\Phi, \bar{\Phi})_*$ is always invariant under the SUSY, then we conclude that the total Wess-Zumino lagrangian is invariant under the 1/2 SUSY transformation and the Lorentz one.

Finally, we summarize main conclusions. We proved some theorems about 1/2 SUSY for N=1 D=4 and N=2 D=2 cases. We discovered the ring $\mathbb{R}[XYZ]$ whose product is defined by the SUSY $*$ product. In other words, $\mathbb{R}[XYZ]$ is closed under the SUSY $*$ product. The SUSY $*$ product does not preserve (anti-)chirality, nevertheless we proved that $\int d^2\theta$ of $\mathbb{R}[XYZ]$ elements are 1/2 SUSY invariant. From these facts, we easily saw that usual F-terms, which take forms of $\int d^2\theta \Phi_*^n$, is invariant under 1/2 SUSY transformation in the framework of the SUSY $*$ deformed theory. Furthermore, $\mathbb{R}[XYZ]$ made the new way to construct 1/2 SUSY invariant observables for both SUSY $*$ formulation and non-SUSY $\star$ formulation. Using this method we can construct explicitly chirality broken observables as 1/2 SUSY invariant observables, for example $\int d\theta^2 D^\alpha * \phi * D_\alpha * \phi * (\phi)_*^n$. Such observables are still 1/2 SUSY observables after replacing $*$ by non-SUSY $\star$ product. In addition, we proved the Lorentz invariance of F-terms and D-terms by explicit calculations. It is possible to show the Lorentz invariance by using ring structure as similar to the proof of 1/2 SUSY.

Acknowledgments

We thank the referee of Phys. Lett. B for many useful suggestions and pointing out typos in earlier version of this article. A.Sako is supported by 21st Century COE Program at Keio University (Integrative Mathematical Sciences: Progress in Mathematics Motivated by Natural and Social Phenomena).

References

- [1] S. Ferrara and M. A. Lledo, JHEP 0005 (2000) 008, hep-th/0002084.
- [2] D. Klemm, S. Penati and L. Tammasia, Class. Quant. Grav. 20 (2003) 2905, hep-th/0104190.
- [3] R. Abbaspur, Int. J. Mod. Phys. A18 (2003) 855, hep-th/0110005.
- [4] R. Abbaspur, hep-th/0206170.
- [5] N. Seiberg, JHEP 0306 (2003) 010, hep-th/0305248.
- [6] S.Ferrara, M. A. Lledo and O. Macia, JHEP 0309 (2003) 068, hep-th/0307039.
- [7] M. Chaichian and A. Kobakhidze, hep-th/0307243.
- [8] I. Bars, C. Deliduman, A. Pasqua and B. Zumino, Phys. Rev. D68 (2003) 106006, hep-th/0308107.
- [9] R. Britto, B. Feng and S.-J. Rey, JHEP 0307 (2003) 067, hep-th/0306215.
- [10] R. Britto, B. Feng and S.-J. Rey, JHEP 0308 (2003) 001, hep-th/0307091.
- [11] O. Lunin and S.-J. Rey, JHEP 0309 (2003) 045, hep-th/0307275.
- [12] D. Berenstein and S.-J. Rey, hep-th/0308049.
- [13] S. Terashima and J.-T. Yee, hep-th/0306237.
- [14] T. Araki, K. Ito and A. Ohtuka, Phys. Lett. B573 (2003) 209, hep-th/0307076.
- [15] R. Britto and B. Feng, Phys. Rev. Lett. 91 (2003) 201601, hep-th/0307165.
- [16] A. Romagnoni, JHEP 0310 (2003) 016, hep-th/0307209.
- [17] W. T. Grisaru, S. Penati and A. Romagnoni, JHEP 0308 (2003) 003, hep-th/0307099.
- [18] M. Alishahiha, A. Ghodsi and N. Sadooghi, hep-th/0309037.
- [19] I. Chepelev and C. Ciocarlie, JHEP 0306 (2003) 031, hep-th/0304118.
- [20] A. Imaanpur, JHEP 0309 (2003) 077, hep-th/0308171.
- [21] R. Abbaspur, hep-th/0308050.
- [22] M. Hatsuda, S. Iso and H. Umetsu, Nucl. Phys. B671 (2003) 217, hep-th/0306251.
- [23] E. Ivanov, O. Lechtenfeld and B. Zupnik, hep-th/0308012.
- [24] S. Ferrara and E. Sokatchev, hep-th/0308021.
- [25] M. Chaichian and D. Polyakov, Phys. Lett. B535 (2002) 321, hep-th/0202209.
- [26] J. de Bohr, P. A. Grassi and P. van Nieuwenhuizen, Phys. Lett. B574 (2003) 98, hep-th/0302078.
- [27] N. Berkovits and N. Seiberg, JHEP 0307 (2003) 010, hep-th/0306226.
- [28] H. Ooguri and C. Vafa, Adv. Theor. Math. Phys. 7 (2003) 53, hep-th/0302109.
- [29] H. Ooguri and C. Vafa, hep-th/0303063.
- [30] H. Kawai, T. Kuroki and T. Morita, Nucl. Phys. B664 (2003) 185, hep-th/0303210.
- [31] J.-H. Park, JHEP 0309 (2003) 046, hep-th/0307060.
- [32] Y. Shibusawa and T. Tada, hep-th/0307236.
- [33] M. R. Douglas and N. A. Nekrasov, Rev. Mod. Phys. 73 (2001) 977, hep-th/0106048.
- [34] J. Wess and J. Bagger, *Supersymmetry and Supergravity*, (Princeton Univ. Press. 1992)